

**3. Majority of species have elongate, conical siphon =**

*Uranotaenia*


*Orthopodomyia*

*Psorophora*

*Culiseta*

*Culex*


*Aedes/Ochlerotatus*




© 2002 Stephen L. Doggett

## Quick key to larvae with “normal” siphons


- 1a. Siphon without pecten.....*Orthopodomyia*
- 1b. Siphon with pecten.....2
- 2a. Head longer than wide,  
comb plate on abdominal segment 8.....*Uranotaenia*
- 2b. Head as wide as long, no comb plate.....3
- 3a. Tufts of setae at base of siphon.....*Culiseta*
- 3b. No basal tufts of setae on siphon.....4
- 4a. Anal segment with complete saddle  
and large ventral brush.....*Psorophora*
- 4b. Not as above.....5
- 5a. Single pair of setae on siphon near midpoint.....*Aedes/Ochlerotatus*
- 5b. Multiple pairs of setae on siphon at midpoint and  
beyond.....*Culex*



Pecten



**NO pecten on siphon  
= *Orthopodomyia***






Figure 24.10





Figure 24.13

Pecten on siphon AND large comb plate with scales on “last” (#8)  
abdominal segment =  
*Uranotaenia*



© 2002 Dept. Medical Entomology, ICPMR



**Pecten AND basal tufts of  
setae on siphon = *Culiseta***





Figure 24.2

**Anal segment with complete “saddle” and obvious ventral brush  
= *Psorophora***

Multiple pairs of setae on siphon =  
*Culex*



Single pair of setae on siphon =  
*Aedes/Ochlerotatus*






Figure 24.29

Single pair of setae on siphon =  
*Aedes/Ochlerotatus*




Multiple pairs of setae on siphon = *Culex*



*Aedes/Ochlerotatus* siphons  
tend to be short/stout  
compared to *Culex*

© 2002 Dept. Medical Entomology, ICPMR

*Culex* siphons tend to be  
long/thin compared to  
*Aedes/Ochlerotatus*



**But note *Culiseta siphon***

